Certificate of Analysis (CoA) for induced Pluripotent Stem Cells



EBISC

This product is for research only

ECACC Catalogue No: 66541182

Cell Line Name	CHDIi035-A	Batch Number	M001
Donor ID	MTM#108		
Disease Association	Huntington's Disease	Phenotype of Donor	Affected
Tissue of Origin	Fibroblasts	Sex	Male
Reprogramming Method	CytoTune™ 2.0 Sendai		
Passage Number	Passage 10	Cell number / vial	1.7 x 10 ⁶
Culture Matrix	Matrigel	Culture Medium	mTESR™1
O ₂ Concentration	21%	CO ₂ Concentration	5%
Passaging Method	EDTA	Additional Culture Information	N/A
Cryopreservation Medium	Cryostor® CS10		
Recommendation for thawing	Recommended to thaw 20% of vial contents into 1 well of a 6-well plate and 80% of vial contents into a second well of a 6-well plate		
	Refer to cell line user protocols for further guidance at www.EBiSC.org		nce at www.EBiSC.org
Additional Comments	Typical recovery after thaw, typical growth to confluency		
Associated Publications	PubMed ID: N/A		

Please see www.EBiSC.org for further information on Quality Control applied to lines released by EBiSC. The following standard testing criteria have been determined within EBiSC, prior to release of this product:

Test	Assay	Acceptance Criteria	Result
Sterility	Inoculation for microbiological growth	Not Detected	Pass
	qPCR for Mycoplasma	Not Detected	Pass
	Virology (HBV, HCV, HIV1, HIV2, HTLV1 & HTLV2)	Not Detected	Pass
Cell Line Identity	Short Tandem Repeat analysis using PCR	N/A	Allele data recorded and available upon request. Profile match to donor fibroblast
Viability	Visual Assessment	Growth to confluence post-thaw	Acceptable



In case of queries, please contact <u>culturecollections.technical@phe.gov.uk</u>. European Collection of Authenticated Cell Cultures (ECACC), Culture Collections, Public Health England, Tel: +44 (0) 1980 612684

Certificate of Analysis (CoA) for induced Pluripotent Stem Cells



This product is for research only

ECACC Catalogue No: 66541182

Test	Assay	Acceptance Criteria	Result
Phenotype	Continuous visual assessment of iPSC colony morphology	Recorded	Typical iPSC colonies with low differentiation levels
Phenotype	Flow Cytometry	SSEA-4 > 70% TRA-1-60 > 70% SSEA-1 < 10% POU5F1 > 70%	Pass
Differentiation Potential	Directed differentiation and qPCR for trilineage markers	Up-regulation of germ layer markers	Endoderm : Detected Mesoderm : Detected Ectoderm : Detected
Karyotype	G-Banding	Normal karyotype	Diploid male karyotype 46, XY
Clearance of Reprogramming Factors	eprogramming qPCR for Sendai backbone		Not detected

The following guidance can be found in the Instructions for Use		
Intended use	Expiry Date	
Product Format	Recommended storage conditions	
Volume	Hazardous Information	

Approved CoA

Signature Come Come

Date 19 Sepan9

