iPS17-00027

Certificate of Analysis

Radboudumc

16-09-2017

Fresh PBMCs nucleofected with episomal vectors OCT3/4, SOX2, KLF4, L-MYC, LIN28
HEP17-00063
PBMC
ID
KCNQ2
3
P6
Essential 8 Flex medium
(cat. no. A2858501, Life Technologies)
Mouse Embryonic Fibroblasts (MEFs)
0.5 mM EDTA
046588; 046591

The three individual clones were analyzed for the following specifications

Test Description	Test method	Test specification	Result
Activation of stem	qPCR	Upregulation of OCT4, SOX2,	Pass
cell markers		NANOG, LIN28 compared with	
		fibroblasts	
Expression of stem	Immunocytochemistry	Expression of Oct4, Nanog,	Pass
cell markers		SSEA-4, Tra1-81	
Mycoplasma	PCR	Negative	Pass
Pluripotency	Differentiation assay	Upregulation of germlayer	N/A*
		specific genes	

*N/A: Not Applicable



Figure 1: Cells prior freezing. A - C, respectively clone 1, 2 and 3 at P6. Scale bar = 1000 μ m.



Activation of stem cell markers

All clones are assessed for activation of stem cell markers before freezing. RNA is isolated and gene expression is assessed by quantitative reverse transcription PCR. Ct values are normalized with the housekeeping gene GUSB (set at 1).



absolute expression, normalize to GusB





relative expression to PBMCs

Figure 3. Pluripotency gene upregulation after reprogramming ($\Delta\Delta$ Ct). The expression fold difference of the iPS clones is relative to the parental PBMCs.

Radboudumc

Expression of stem cell markers

Undifferentiated iPS cell clones are stained for the nuclear markers NANOG and OCT4 and surface antigens SSEA4 and TRA1-81. All markers are expressed in pluripotent human stem cells.



B. iPS17-00027 clone 2 P7



Radboudum



C. iPS17-00027 clone 3 P6

Figure 4. Immunofluorescence staining of the iPS cell clones with pluripotency markers.

- 🖉 Pass
- Fail
- Other:

Sev. Abu

Silvia Albert, PhD Supervisor, Radboud Stem Cell Technology Center Date

Frans Cremers, PhD

Head, Radboud Stem Cell Technology Center 2 · (o. 2017 Date