

**Certificate of Analysis 2018**

Invoice number: SCTC2018-00051

Name investigator: Nael Nadif Kasri.....

Cell line number: IPS18-00047.....

Project name: KCNQ2 encefalopathie.....


**Table 1: Information reprogrammed cell line**

Information cell line:	
Product description	Fresh PBMCs nucleofected with vectors containing the genes OCT3/4, SOX2, KLF4, L-MYC, LIN28
Parental cell line	HEP18-00105
Parental cell type	PBMC
Diagnosis	ID
Mutation	N/A*
Number of clones	3
Passage (P) of iPS cells reported at submission	P6
Culture medium	Essential 8 Flex medium
Culture matrix	Vitronectin
Feeder during reprogramming	Mouse Embryonic Fibroblasts (MEFs)
Passage method	0.5 mM EDTA
Protocols in Q-portal	046588; 046591

**Table 2: Information characterization reprogrammed cell line**

Test Description:	Test method:	Test specification:	Result:
Activation of stem cell markers	qPCR	Upregulation of <i>OCT4</i> , <i>SOX2</i> , <i>DNMT3b</i> , <i>LIN28</i> compared to PBMCs	Pass
Expression of stem cell markers	Immunocytochemistry	Expression of OCT4, NANOG, SSEA4, TRA-1-81	Pass
Mycoplasma	PCR	Negative	Pass
Pluripotency	Differentiation assay	Upregulation of germlayer specific genes	N/A*

\*N/A: Not Applicable



**Figure 1: Cells prior to freezing. A - C, respectively clone 1 and 2 at P6. Scale bar = 1000 µm. Picture of clone 3 not available.**

Activation of stem cell markers

All clones are assessed for activation of stem cell markers before freezing. RNA is isolated and gene expression is assessed by quantitative reverse transcription PCR. Ct values are normalized with the housekeeping gene GUSB (set at 1).

Absolute expression, normalized to GusB




Figure 2: Gene expression of three iPS cell clones compared to the parental PBMCs ( $\Delta$ Ct).

Expression relative to parental PBMCs






Figure 3: Pluripotency gene upregulation after reprogramming ( $\Delta\Delta$ Ct). The expression fold difference of the iPS clones is relative to the parental PBMCs.

## Expression of stem cell markers

Undifferentiated iPS cell clones are stained for the nuclear markers NANOG and OCT4 and surface antigens SSEA4 and TRA-1-81. All markers are expressed in pluripotent human stem cells.

**A. *iPS18-00047 clone 1 P6*****B. *iPS18-00047 clone 2 P6***

C. iPS18-00047 clone 3 P6




Figure 4. Immunofluorescence staining of the iPS cell clones with pluripotency markers.

- Pass
- Fail
- Other:

*Silvia Albert*

---

**Silvia Albert, PhD**  
 Manager, Radboud Stem Cell Technology Center  
 Date 21.09.2018