

Contents

1. Description

1.1 Background information

1.2 Applications

1.3 Reagent requirements

2. Protocol

2.1 Preparation of complete media

2.2 Passaging of human ES or iPS cells

1. Description

Components 500 mL StemMACS™ iPS-Brew XF, Basal Medium

10 mL StemMACS iPS-Brew XF, 50× Supplement

Specifications pH: 7.2–7.6

Osmolality: 300–340 mOsmol/kg

Quality control Maintenance of human iPS cell morphology and surface phenotype (TRA-1-60⁺, SSEA-4⁺) over five continuous passages. Low endotoxin level by Limulus Amoebocyte Lysate (LAL) assay. Tested negative for mycoplasma.

Storage Store the StemMACS iPS-Brew XF, Basal Medium protected from light at 2–8 °C. Do not freeze. The expiration date is indicated on the vial label.

Upon arrival store StemMACS iPS-Brew XF, 50× Supplement at –20 °C.

Aliquots of the supplemented complete media can be stored at –20 °C for up to 2 month. Avoid repeated freeze-thaw-cycles. Once thawed, aliquots should be kept at 2–8 °C and be used within 2 weeks. The expiration date is indicated on the vial label.

Intended use

StemMACS iPS-Brew XF is intended for research use. It is not intended for human or animal diagnostic or therapeutic use.

1.1 Background information

StemMACS iPS-Brew XF is a xeno-free media formulation for the maintenance and expansion of human pluripotent stem cells under feeder-free conditions. StemMACS iPS-Brew XF supports rapid adaption of feeder-based cultures to a feeder-free environment. The formulation is compatible with standard cell attachment matrices, e.g. Matrigel® or vitronectin. It enables robust and efficient expansion of human embryonic stem cells (ES) or induced pluripotent stem cells (iPS) over multiple passages while maintaining a pluripotent phenotype as well as pluripotent differentiation potential. StemMACS iPS-Brew XF allows rapid culture re-initiation of pluripotent stem cell cultures after cryopreservation.

1.2 Applications

- Culture of human ES or iPS cells under xeno- and feeder-free conditions
- Rapid and easy adaption of feeder-based culture to a feeder-free environment
- Rapid culture initiation after cryopreservation

1.3 Reagent requirements

- Buffer: Dulbecco's phosphate-buffered saline (DPBS) without Ca²⁺ and Mg²⁺.
- A small molecule ROCK inhibitor, e.g., StemMACS Y27632 (# 130-103-922) or StemMACS Thiazovivin (# 130-104-461) to improve cell attachment and survival.
- StemMACS Passaging Solution XF (# 130-104-688) for passaging in cell clusters.
- 0.05% Trypsin/EDTA (alternatively, Accutase® or TrypLE™) and Soybean Trypsin Inhibitor (0.5 mg/mL) for single cell splitting.
- Cell attachment substrate. Validated substrates are, e.g., Matrigel®, Geltrex®, Laminin-511, Lamin-521, iMatrix-511, vitronectin, or CTS™ CELLstart™ substrate
- 15 mL conical tubes

2. Protocol

2.1 Preparation of complete media

Before StemMACS iPS-Brew XF can be used in cell culture, the two kit components need to be mixed according to the following protocol to obtain the complete medium.

1. Thaw StemMACS iPS-Brew XF, 50× Supplement at 2–8 °C prior to use.
2. To obtain the complete medium add 10 mL StemMACS iPS-Brew XF, 50× Supplement to 500 mL StemMACS iPS-Brew XF, Basal Medium. Mix well. The media is ready-to-use now. Use the complete medium within 2 weeks when stored at 2–8 °C.
3. For longer storage, prepare 50 mL aliquots and store at –20 °C for up to 2 month. Thaw aliquots of complete medium overnight at 2–8 °C. Once thawed, keep aliquots at 2–8 °C and use within 2 weeks.

2.2 Passaging of human ES or iPS cells

▲ StemMACS iPS-Brew XF is compatible with standard passaging techniques such as traditional colony cutting, passaging in cell clusters or single cells. It is recommended to use the single-cell or cluster-splitting technique in the presence of a small molecule ROCK inhibitor.

Protocol for passaging in cell clusters

1. Coat 6-well plates with an appropriate attachment substrate according to the manufacturer's instructions.

2. Aspirate the cell culture supernatant.
3. Wash the cell layer with 3 mL of buffer per well.
4. Add 1 mL of StemMACS Passaging Solution XF per well. Gently rock the plate to distribute the solution evenly.
5. Incubate at room temperature for 4 minutes. Monitor the detachment process under the microscope.

▲ **Note:** Colonies must not detach completely. Only wait until the colony edges lift off (see figure 2).




Figure 1: Colonies before addition of StemMACS Passaging Solution XF.




Figure 2: After 4 minutes incubation, colony edges start to lift off. At this point, the passaging solution should be removed.

6. Carefully remove the StemMACS Passaging Solution XF.
7. Per well, add 3 mL of StemMACS iPS-Brew XF supplemented with ROCK inhibitor (e.g., 2 μ M StemMACS Thiazovivin or 10 μ M StemMACS Y27632).
8. Gently detach the colonies by rinsing the well with a 5 mL serological pipette.
9. Transfer the cell suspension into a 15 mL conical tube.
10. Carefully pipette up and down 2–3 times to break up the colonies into smaller cell clusters.

▲ **Note:** Take care to minimize break-up of colonies. Do not create single cells!
11. Transfer the cell clusters into a fresh, appropriately coated 6-well cell culture plate. Use 2 mL StemMACS iPS-Brew XF supplemented with ROCK inhibitor per well and a splitting ratio between 1:6 and 1:20.

▲ **Note:** The optimal splitting ratio will depend on the cell line and must be determined empirically.

12. After 48 hours, replace media with fresh StemMACS iPS-Brew XF without ROCK inhibitor and continue with daily media changes.

▲ **Note:** Many ES and iPS cell lines will also tolerate every-other-day media changes when using StemMACS iPS-Brew XF.

Protocol for single-cell splitting

1. Coat 6-well plates with an appropriate attachment substrate according to the manufacturer's instructions.
2. Aspirate cell medium, wash each well with 3 mL of buffer.
3. Add 0.7 mL of 0.05% Trypsin/EDTA per well (alternatively, use Accutase or TrypLE). Gently rock the plate to ensure even distribution of the enzyme solution.
4. Incubate for 5 minutes at 37 °C.
5. Stop enzymatic reaction by adding 2 mL of Soybean Trypsin Inhibitor (0.5 mg/mL) per well.
6. Using a 5 mL serological pipette, dissociate to a single-cell suspension by carefully pipetting up and down.
7. Determine cell number.
8. Depending on the cell line, seed 70,000–150,000 cells per well (7000–16,000 cells/cm²). Transfer the desired cell number into a 15 mL conical tube.
9. Centrifuge for 5 minutes at 200 \times g.
10. Aspirate supernatant.
11. Resuspend the cell pellet in StemMACS iPS-Brew XF supplemented with a small molecule ROCK inhibitor (10 μ M StemMACS Y27632 or 2 μ M StemMACS Thiazovivin). Use 2 mL medium per well.
12. After 48 hours, replace media with fresh StemMACS iPS-Brew XF without ROCK inhibitor and continue with daily media changes.

▲ **Note:** Many ES and iPS cell lines will also tolerate every-other-day media changes when using StemMACS iPS-Brew XF.

All protocols and data sheets are available at www.miltenyibiotec.com.

Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

Accutase is a registered trademark of Innovative Cell Technologies, Inc.

Matrigel is a registered trademark of Corning Incorporated.

CTS, CELLstart, Geltrex, and TrypLE are trademarks of Life Technologies Corporation.

MACS is a registered trademark and StemMACS is a trademark of Miltenyi Biotec GmbH.

Copyright © 2015 Miltenyi Biotec GmbH. All rights reserved.